

CORRECTION

Open Access



Correction: Amniotic fluid-derived exosomes attenuated fibrotic changes in POI rats through modulation of the TGF- β /Smads signaling pathway

Nahideh Nazdikbin Yamchi¹, Shahin Ahmadian¹, Halimeh Mobarak¹, Farhad Amjadi¹, Rahim Beheshti², Amin Tamadon^{3,4}, Reza Rahbarghazi^{5,6} and Mahdi Mahdipour^{1,7*}

Correction: *J Ovarian Res* 16, 118 (2023)

<https://doi.org/10.1186/s13048-023-01214-1>

Following publication of the original article [1], the authors reported that Figs. 3 and 4 have been updated due to unintentional errors when compiling the data. The correct figures were included in this correction.

The online version of the original article can be found at <https://doi.org/10.1186/s13048-023-01214-1>.

*Correspondence:

Mahdi Mahdipour

mahdi.mahdipour@gmail.com; mahdipourm@tbzmed.ac.ir

¹Stem Cell Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

²Faculty of Veterinary Medicine, Shabestar Islamic Azad University, Shabestar, Iran

³PerciaVista R&D Co., Shiraz, Iran

⁴Department for Scientific Work, Marat Ospanov Medical University, West, Aktobe, Kazakhstan

⁵Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran


⁶Department of Applied Cell Sciences, Faculty of Advanced Medical Sciences, Tabriz University of Medical Sciences, Tabriz, Iran

⁷Department of Reproductive Biology, Faculty of Advanced Medical Sciences, Tabriz University of Medical Sciences, Tabriz, Iran




© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Corrected figure 3:



Corrected figure 4:



The original article [1] has been corrected.

Published online: 19 August 2024

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

1. Nazdikbin Yamchi N, Ahmadian S, Mobarak H, et al. Amniotic fluid-derived exosomes attenuated fibrotic changes in POI rats through modulation of the TGF- β /Smads signaling pathway. *J Ovarian Res.* 2023;16:118. <https://doi.org/10.1186/s13048-023-01214-1>.